

SERANDANG

Ocellated Snakehead

Channa pleurophthalma

Animalia
Chordata
Actinopterygii
Perciformes
Channidae
Channa
Channa pleurophthalma



Local name: *Kerandang, toman bunga, selendang mayang.*



Habitat

Living in rivers, floodplain, and swamps and in areas connected to rivers which when flooded will be inundated and fish will live in it.



Distribution

This fish is found in Indonesia, with the distribution area in Sumatra and Kalimantan.



Food

This fish is a type of carnivorous fish which preys on insects, small fish, and shrimp as food.



Reproduction

It spawns throughout the year with peak spawning period at the beginning of the rainy season. The fecundity of serandang fish ranges from 1,223-4,290 eggs.

SERANDANG

Ocellated Snakehead

Channa pleurophthalma

Animalia
Chordata
Actinopterygii
Perciformes
Channidae
Channa
Channa pleurophthalma



Kerandang, toman bunga, selendang mayang.



Habitat

Hidup di perairan sungai, rawa banjiran, dan lebak. Daerah yang terhubung dengan sungai yang apabila banjir akan tergenang dan ikan akan hidup di dalamnya.



Sebaran

Keberadaan ikan ini tercatat ditemukan Indonesia, dengan wilayah sebaran adalah di Sumatera dan Kalimantan.



Makanan

Ikan ini termasuk jenis ikan karnivora dengan makanan berupa serangga, ikan berukuran kecil, dan udang,



Reproduksi

Memiliki pola pemijahan sepanjang tahun dengan puncak memijah pada awal musim penghujan. Fekunditas ikan serandang berkisar 1.223 - 4.290 butir telur.