

SEMURINGAN

Striped Barb

Puntius sp.

Animalia
Chordata
Actinopterygii
Cypriniformes
Cyprinidae
Puntius
Puntius sp.



Local name: *Semuringan*.



Habitat

Living in rivers, peatland, acidic waters, and slow water flow with a pH of 3-6.5, an aquatic environment where there are a lot of submerged grass or aquatic plants.



Distribution

These fish are found in Indonesia, Malaysia, and Thailand. In Indonesia, the distribution area is in Sumatra, Bangka, and Kalimantan.



Food

It's a type of omnivorous fish species which preys on zooplankton, insect larvae, crustaceans, worms, organic detritus, and aquatic plants.



Reproduction

It has a spawning pattern throughout the year with peak spawning period at the beginning of the rainy season. The fecundity of serandang fish ranges from 1,000 to 2,275 eggs.

SEMURINGAN

Striped Barb

Puntius sp.

Animalia
Chordata
Actinopterygii
Cypriniformes
Cyprinidae
Puntius
Puntius sp.

Semuringan.




Habitat

Hidup di perairan sungai, rawa gambut, perairan asam, dan perairan yang tenang dengan pH 3 - 6,5. Lingkungan perairan yang banyak terdapat rumput atau tanaman air yang terendam.



Sebaran

Keberadaan ikan ini tercatat ditemukan di Indonesia, Malaysia, dan Thailand. Di Indonesia sendiri wilayah sebarannya adalah di Sumatera, Bangka, dan Kalimantan.



Makanan

Ikan ini termasuk jenis ikan omnivora dengan makanan berupa zooplankton, larva serangga, krustasea, cacing, detritus organik dan tanaman air.



Reproduksi

Memiliki pola pemijahan sepanjang tahun dengan puncak memijah pada awal musim penghujan. Fekunditas ikan serandang berkisar 1.000 - 2.275 butir telur.