



SEPAT SIAM

Snakeskin Gourami

Trichogaster pectoralis

Animalia
Chordata
Actinopterygii
Anabantiformes
Anabantoidei
Osphronemidae
Trichogaster
Trichogaster pectoralis

Local name: Sepat siam, sepat, sliper (East Java).



Habitat

The fish lives in fresh water, benthopelagic with calm, shallow currents, and lots of aquatic plants, swamps, lakes, ditches, rice fields, and slow water flow. It can live in waters with pH ranging from 4-9 and the temperature is 22°C-28°C.



Distribution

This fish is naturally (natively) found in Cambodia, Laos, Viet Nam, and Thailand. It is an introduced fish in Indonesia, Malaysia, Myanmar, Philippines, Singapore, Sri Lanka, New Caledonia, Colombia, and Papua New Guinea. The distribution area in Indonesia is in Sumatra, Sulawesi, Kalimantan, Papua, and Java.



Food

It is included in type of omnivorous fish whose food includes mosses, aquatic plants, small animals that fit the size of their mouths.



Reproduction

It begins to breed at the age of 6 - 8 months, with a weight ranging from 95 - 120 g and a total length of more than 8 - 21 cm. Spawning occurs in the rainy season and in nests in the form of bubbles (foam) that have been prepared beforehand.

SEPAT SIAM

Snakeskin Gourami

Trichogaster pectoralis

Animalia
Chordata
Actinopterygii
Anabantiformes
Anabantoidei
Osphronemidae
Trichogaster
Trichogaster pectoralis

Sepat siam, sepat, sliper (Jawa Timur).



Habitat

Hidup pada perairan air tawar, benthopelagik dengan arus yang tenang, dangkal, dan banyak tanaman air. Kisaran pH air 4 - 9 dan suhu 22°C - 28°C. Rawa-rawa, danau, parit, sawah dan sungai yang berair tenang.



Sebaran

Keberadaan ikan ini secara alami (asli) adalah di Kamboja, Laos, Vietnam, dan Thailand. Sebagai ikan introduksi di Indonesia, Malaysia, Myanmar, Phillipina, Singapura, Sri Lanka, New Caledonia, Kolombia, dan Papua New Guinea. Wilayah sebaran di Indonesia adalah di Sumatera, Sulawesi, Kalimantan, dan Jawa.



Makanan

Termasuk jenis ikan omnivora, mulai dari lumut, tanaman air, hewan kecil yang muat dengan ukuran mulutnya.



Reproduksi

Mulai berbiak pada umur 6 - 8 bulan, dengan bobot berkisar 95 - 120 gr dan panjang total lebih dari 8 - 21 cm. Memijah pada musim penghujan dan terjadi pada sarang yang berupa gelembung (busa) yang telah disiapkan sebelumnya.